



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>5</sup> :</b> C07K 15/06, 15/14, C12N 15/04 C12N 15/06, 15/10, 15/12	<b>A1</b>	<b>(11) International Publication Number:</b> WO 93/22348 <b>(43) International Publication Date:</b> 11 November 1993 (11.11.93)
<b>(21) International Application Number:</b> PCT/US93/03614 <b>(22) International Filing Date:</b> 16 April 1993 (16.04.93) <b>(30) Priority data:</b> 07/872,304 24 April 1992 (24.04.92) US <b>(71) Applicant:</b> BAYLOR COLLEGE OF MEDICINE (US/ US); One Baylor Plaza, Houston, TX 77030 (US). <b>(72) Inventors:</b> CONNEELY, Orla, M. ; 2601 South Braes- wood, #1307, Houston, TX 77025 (US). HEADON, De- nis, R. ; Lishin, Aghnacorra, Dangan Lower, Galway (IE). O'MALLEY, Bert, W. ; 639 Ramblewood, Hous- ton, TX 77079 (US). MAY, Gregory, S. ; 4119 Durness, Houston, TX 77025 (US).		<b>(74) Agent:</b> PAUL, Thomas, D.; Fulbright & Jaworski, 1301 McKinney, Suite 5100, Houston, TX 77010-3095 (US). <b>(81) Designated States:</b> AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, I.K., LU, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, SK, Eu- ropean patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> PRODUCTION OF RECOMBINANT HUMAN LACTOFERRIN  <b>(57) Abstract</b> <p>The present invention provides novel plasmids, transfected eucaryotic cells and methods of producing these plasmids and transfected eucaryotic cells. The novel plasmid contains the cDNA for human lactoferrin protein. Methods for the production of human lactoferrin protein in <i>A. Oryzae</i> are also provided. Thus, the present invention provides an efficient and economical means for the production of recombinant human lactoferrin protein.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SK	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LI	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LJ	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VB	Viet Nam
FI	Finland				

-1-

## PRODUCTION OF RECOMBINANT HUMAN LACTOFERRIN

### BACKGROUND OF THE INVENTION

#### Field of the Invention

5           The present invention relates generally to the field of iron-binding glycoproteins. More specifically, the present invention relates to the recombinant production of human lactoferrin.

#### Description of the Related Art

10           Human lactoferrin (LF) is a member of the transferrin family of iron-binding monomeric glycoproteins. It was originally discovered in milk where it can reach levels of 7 grams/liter in colostrum. LF has since been detected in other external fluids such as tears, saliva and mucosal secretions and also in the secondary granules of polymorphonuclear leukocytes.

15           LF is a 78 kDa glycoprotein having a bilobal structure with a high degree of homology between the C and N terminal halves which is evident at both the amino acid and three dimensional structural level. Each of these lobes can reversibly bind one ferric iron with high affinity and with the concomitant binding of bicarbonate. The biological functions  
20           proposed for lactoferrin include protection against microbial infection,

-2-

enhanced intestinal iron absorption in infants, promotion of cell growth, regulation of myelopoiesis and modulation of inflammatory responses.

Filamentous fungi have been successfully employed as hosts in the industrial production of extracellular glycoproteins. Certain industrial strains are capable of secreting gram quantities of these proteins. In addition, filamentous fungi are able to correctly perform post-translational modifications of eucaryotic proteins and many strains have U.S Food and Drug Administration approval. Furthermore, large scale fermentation technology and downstream processing experience is available.

Currently, there is no efficient and economical way to produce human LF. Consequently, a long felt need and description in this art would be met by the development of an efficient method for the production of human lactoferrin for nutritional and therapeutic applications and for further investigation into its mechanism of action.

#### SUMMARY OF THE INVENTION

In one embodiment, the present invention provides for a recombinant plasmid comprising the cDNA of human lactoferrin. The plasmid of the present invention is adapted for expression in an eucaryotic cell and contains the regulatory elements necessary for the expression of the human lactoferrin cDNA in this eucaryotic cell.

In another embodiment, the present invention provides for a transformed eucaryotic cell which includes a recombinant plasmid. The eucaryotic cell is selected from a group of filamentous fungi comprising *Aspergillus*. The plasmid contains a plasmid vector into which a polydeoxyribonucleotide segment coding for human lactoferrin protein has been inserted.

In yet another embodiment of the present invention, there is provided a process for producing recombinant human lactoferrin which comprises culturing a transformant eucaryotic cell, which includes a

-3-

recombinant plasmid. The plasmid contains a plasmid vector having a polydeoxyribonucleotide coding for the human lactoferrin protein. After culturing in a suitable nutrient medium until human lactoferrin protein is formed, the human lactoferrin protein is isolated.

5 In still yet another embodiment of the present invention, there is provided a recombinant expression vector. This vector comprises a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression; (2) cDNA coding for human lactoferrin; (3) appropriate transcription and translation initiation  
10 and termination sequences; and (4) a genetic element for selection of aspergillus spores that have been transformed with the vector.

In still yet another embodiment of the present invention, there is provided a method for producing biologically active recombinant lactoferrin. The method comprises synthesizing sequences containing a  
15 selectable marker gene, a promotor, a transcription termination sequence, and a linker sequence; cloning the sequences to form a plasmid; digesting the plasmid with a restriction endonuclease; inserting a cDNA coding for lactoferrin into a restriction site; and transforming eucaryotic cells with the plasmid expressing lactoferrin cDNA.

#### 20 BRIEF DESCRIPTION OF THE DRAWINGS

So that the manner in which the above recited features, advantages, and objects of the invention, as well as others which will become clear, are obtained and can be understood in detail, more particular descriptions of the invention briefly summarized above may be  
25 had by reference to certain embodiments thereof which are illustrated in the appended drawings. These drawings form a part of this specification. It is to be noted, however, that the appended drawings illustrate preferred embodiments of the invention and therefore not to be considered limiting of its scope. The invention may admit to other equally effective equivalent  
30 embodiments.

-4-

Fig. 1 depicts a schematic representation of the *aspergillus oryzae* expression plasmid, pAhlfg.

Fig. 2 shows a southern blot analysis of transformed *aspergillus oryzae* strains.

5 Fig. 3 depicts an RNA analysis of transformant versus control A07.

Fig. 4 shows the silver stained SDS-acrylimide gel analysis of recombinant LF secretion and purification.

10 Fig. 5 illustrates the characterization of recombinant human LF.

Fig. 6 depicts the cDNA sequence for human lactoferrin.

## DETAILED DESCRIPTION OF THE INVENTION

### DEFINITIONS

15 For the purposes of the present application, the term "transferrin family" means a family of iron transferring proteins including serum transferrin, ovotransferrin and lactoferrin. These proteins are all structurally related.

20 For the purposes of the present application, the term "vector(s)" means plasmid vehicle to allow insertion, propagation and expression of lactoferrin cDNA.

For the purposes of the present application, the term "host(s)" means any eucaryotic cell that will allow integration of the lactoferrin expression plasmid into its genome.

25 For the purposes of the present application, the term "promotor(s)" means regulatory DNA sequences that controls transcription of the lactoferrin cDNA.

30 For the purposes of the present application, the term "multiple cloning cassette" means a DNA fragment containing restriction enzyme cleavage sites for a variety of enzymes allowing insertion of a variety of cDNAs.

-5-

For the purposes of the present application, the term "transformation" means uptake of plasmid by a relevant eucaryotic cell.

For the purposes of the present application, the term "iron binding capacity" means ability to bind  $^{59}\text{Fe}$ . Fully functional lactoferrin can bind two atoms of iron per molecule of LF.

For the purposes of the present application, the term "biological activity/biological active" means biological activity of lactoferrin as measured by its ability to bind iron. The lactoferrin protein functions as an iron transfer protein and must bind iron to be biologically active.

All literature references cited in this specification are hereby expressly incorporated by reference.

The following examples are given for the purposes of illustrating various embodiments of the present invention and are not meant to be limitations of the present invention in any form.

### Example 1

#### Fungal strains and transformation

The *pyrG* mutant strain used in these studies was derived from *A. oryzae* (A07 11488). The *pyrG* gene from *A. oryzae* was mutated with 4-nitroquinoline-1-oxide. The *Aspergillus* transformation was carried out by a modification of the procedure of Osmani, et al., *J. Cell. Biol.* 104:1495-1504 (1987). Conidia ( $1 \times 10^6/\text{ml}$ ) were inoculated into 50 ml of YG medium (0.5% yeast extract 2% glucose) containing 5 mM uracil and 10 mM uridine. Growth was at 32°C for 14-16 hours until a germ tube was visible. The germinated conidia were harvested by centrifugation and resuspended in 40 ml of lytic mix containing 0.4 M ammonium sulphate, 50 mM potassium citrate (pH 6.0), 0.5% yeast extract, 0.12 g novozyme, 0.1g Driselase, 100  $\mu\text{l}$   $\beta$ -glucuronidase, 0.5% sucrose and 10 mM  $\text{MgSO}_4$ . Protoplasting was for 2-3 hours at 32°C and 150 rpm. Following protoplasting, filtration using sterile miracloth was necessary to remove

-6-

any undigested mycelia. The protoplasts were harvested by centrifugation and washed twice with 10 ml of 0.4 M ammonium sulphate, 1% sucrose and 50 mM potassium citrate (pH 6.0) at 4°C, resuspended in 1 ml of 0.6 M KCl; 50 mM CaCl<sub>2</sub>; 10 mM Tris-HCl (pH 7.5) and placed on ice. The transformation was performed immediately following the protoplast preparation. Aliquots (100 µl) of the protoplast were added to 3 µg of DNA and 50 µl of 40% polyethylene glycol (PEG) 6000, 50 mM CaCl<sub>2</sub>, 0.6 M KCl and 10 mM Tris-HCl (pH 7.5). The samples were incubated on ice for fifteen minutes after which an additional 1 ml of the PEG solution was added and incubation at room temperature was continued for thirty minutes. Aliquots of this mixture were plated in 3 ml of 0.7% minimal media, supplemented with 0.4% ammonium sulphate onto plates containing the same but solidified with 2% agar. All subsequent growth was at 32°C.

## Example 2

### Plasmid Construction

A schematic representation of the expression plasmid is shown in Fig. 1. The complete cDNA encoding human LF was repaired using the Klenow fragment of DNA polymerase I and subcloned into AccI digested and repaired pGEM4 to generate pGEMhLFc. In order to remove the LF signal sequence and generate a 5' end in frame with the  $\alpha$ -amylase sequences, a 252 base pair lactoferrin fragment (nt 69-321) containing HindII/AccI ends was obtained by polymerase chain reaction (PCR) amplification of pGEMhLFc plasmid DNA. The oligo primers used were as follows: the 5' end oligonucleotide as shown in SEQ. ID. No. 1:

(CTGGGTCGACGTAGGAGAAGGAGTGTTCAGTGGTGC)

and the 3' end oligonucleotide as shown in SEQ. ID. No. 2:



-7-

(GCCGTAGACTTCCGCGCTACAGG).

This PCR fragment was digested with HindII and AccI and was subcloned into Hind II/AccI digested pGEMhLFC generating pGEMhLF. A 681 base pair  $\alpha$ -amylase fragment with Asp718/PvuII ends encoding the promotor, signal sequence and the alanine residue from the start of the mature  $\alpha$ -amylase II gene, was obtained by PCR amplification of *A. oryzae* genomic DNA. The oligo primers were as follows: the 5' end oligonucleotide as shown in SEQ. ID. No. 3:

(GAGGTACCGAATTCATGGTGTTTTGATCATTTTAAATTTTATAT)

and the 3' end oligonucleotide as shown in SEQ. ID. No. 4:

(AGCAGCTGCAGCCAAAGCAGGTGCCGCGACCTGAAGGCCGTACAG).

The amplified DNA was digested with Asp718 and PvuII and subcloned into Asp718/HindII digested pGEMhLF. The resulting plasmid (pGEMAhLF) was digested with EcoRI and the resulting 2.8 kb  $\alpha$ -amylase-lactoferrin fragment was subcloned into a unique EcoRI site in pAL3 according to the method of generating pAhLF\*. Synthetic oligonucleotide were used to provide the last five carboxy terminal codons of lactoferrin (nt 2138 - 2153) missing in pAhLF\* and also to provide the first 180 bp of 3' untranslated sequences from the *A. niger* glucoamylase gene. The resulting plasmid (pAhLFG) was used to transform the *A. oryzae* pyrG mutant strain.

With reference to FIGURE 1, *Aspergillus oryzae* expression plasmid, pAhLFG contains 681 bp of 5'-flanking sequence of the *A. oryzae* AMY11 gene which includes the signal sequence and first codon of mature  $\alpha$ -amylase. The cDNA coding for mature human lactoferrin is subcloned in frame downstream from these sequences allowing recombinant protein

-8-

production by the addition of starch to the growth medium. The *Aspergillus niger* glucoamylase 3' untranslated region provides the transcription terminator and polyadenylation signals. The plasmid also contains the *Neurospora crassa* pyr4 selectable marker and an ampicillin resistance gene.

The plasmid construct (pAhLFG) used for expression of human LF contains a 681 bp fragment that encodes the promotor and secretory signal peptide of the *A. oryzae*  $\alpha$ -amylase II gene (AMY11). The signal sequence also contains the codon for alanine from the start of the  $\alpha$ -amylase mature protein generating the signal sequence cleavage site (Leu Ala Ala) recognizable by an endogenase  $\alpha$ -amylase peptidase. A human lactoferrin cDNA fragment encoding the mature protein was subcloned in frame immediately downstream from the AMYII sequences, placing it under the control of this highly efficient starch inducible promoter. In order to stabilize the transcribed human LF mRNA, a 180 bp fragment encoding the 3' untranslated region of the glucoamylase gene from *Aspergillus niger* was ligated into a unique BamHI site in the multiple cloning cassette, immediately downstream of the human LF cDNA providing the transcription terminator and polyadenylation signals. The plasmid also contains the *Neurospora crassa* Pyr4 selectable marker which complements a pyrG auxotrophic mutation of *A. oryzae* and allows for selection of spores that have been transformed with the plasmid by growth in the absence of uridine.

### Example 3

#### Genomic DNA Manipulation

*A. oryzae* DNA was isolated from 200 mg of lyophilized mycelia as described by Rafnussen, et al., *J. Biol. Chem.*, 265:13767-13775 (1990). The DNA was digested with EcoRI, size fractionated on a 0.8% agarose gel and transferred to nitrocellulose. Prehybridization and hybridization of the nitrocellulose filter for Southern analysis were

-9-

performed in 6XSSC, 0.1% SDS and 0.5% dried milk at 65°C for 16 hours. Hybridization solution contained  $1 \times 10^7$  cpm  $^{32}$ P-labelled lactoferrin cDNA probe (2.1 Kb). The filter was washed in 2XSSC, 0.5% SDS at room temperature for 30 minutes followed by two washes in 0.5X SSC, 0.5% SDS at 68°C for 30 minutes. The filter was dried, exposed at -70°C for two hours and developed by autoradiography.

With reference to FIGURE 2, Southern blot analysis was performed on transformed *Aspergillus oryzae* strains. Genomic DNA from individual transformants and control AO7 were hybridized with a radiolabelled hLF cDNA probe (2.1 kb). The arrow points to a radiolabelled fragment (2.8 kb) generated upon EcoRI digestion of the expression plasmid which is present in all the transformants (#1-9) but is absent in control untransformed AO7. Molecular weights of bacteriophage lambda Hind 111 fragments are indicated at the left.

#### Example 4

##### Northern Analysis

RNA was isolated from lyophilized mycelia (200 mg) using commercially available RNazol B (Biotecx Laboratories, INC, Houston, TX) according to the manufacturers instructions. Total RNA (20 µg) was electrophoresed in a 0.8% agarose gel containing 2.2 M formaldehyde. The RNA was transferred to nitrocellulose and hybridized with either a 2.1 kb lactoferrin cDNA or a 1.8 kb genomic  $\alpha$ -amylase fragment corresponding to the coding region of the  $\alpha$ -amylase 11 gene. The probes were  $^{32}$ P-labelled by nick translation (specific activity  $2 \times 10^6$  cpm/ug). Hybridization was carried out 2 x SSC, .05% dried milk at 65°C over an ice with  $2 \times 10^6$  cpm probe/ml.

Washes were identical to those employed in the Southern analysis. The filters were dried, exposed at -70°C for two hours and developed by autoradiography. RNA dot blots were performed using nitrocellulose membrane and the manifold dot blot system. Hybridization

-10-

and washing conditions were as described above for Southern analysis. Radioactivity was quantitated using the betagon blot analyzer.

Recombinant production of lactoferrin protein has been described in its preferred embodiment. However, it could also be produced in a number of other sources such as fungal sources such as *saccharomyces cerevisiae* or *pichia pastorsis* or insect cells such as SF9.

With reference to FIGURE 3, RNA analysis of transformant versus control AO7 was performed. In Panel A, Northern analysis of RNA (20  $\mu$ g) from control AO7 and transformant #1 hybridized with radiolabelled human LF cDNA. Human LF mRNA (2.3 kb) was detected in the transformant #1 but not in the control untransformed AO7. The positions of the 28s and 18s rRNA bands are indicated on the left. In Panel B, Dot blots of RNA (5 and 10  $\mu$ g) from control AO7 versus transformant #1 using a radiolabelled  $\alpha$ -amylase genomic DNA probe. In Panel C, Dot blots of RNA (5 and 10  $\mu$ g from control AO7 and transformant #1 using radiolabelled human LF cDNA probe as illustrated.

Northern analysis was performed to determine if lactoferrin mRNA was transcribed correctly and efficiently in *A. oryzae* under the regulatory control elements of our expression plasmid. Spores ( $1 \times 10^6$ /ml) from transformant #1 and from control untransformed spores were inoculated into fungal medium containing 1.5% glucose as carbon source and grown at 30°C for 48 hours in small shake flask cultures. The cultures were washed and reinoculated into fungal medium containing 3% starch to induce transcription of the human LF mRNA. After 24 hours, the cells were harvested and RNA was isolated. Total RNA (20  $\mu$ g) was size fractionated on a 1.0% agarose gel containing 2.2 M formaldehyde and blotted on nitrocellulose.

Human lactoferrin mRNA was detected using  $^{32}$ P labelled human LF cDNA (2.0 kb) probe. Hybridization with human LF radiolabelled cDNA probe detected a specific radiolabelled band at the correct size for lactoferrin mRNA (2.3kb) in the transformant but not in

-11-

the control untransformed strain (Fig. 3A). Quantitation of mRNA levels by dot assay showed comparable levels of expression of endogenous  $\alpha$ -amylase rRNA between control AO7 and transformant #1 (Fig. 3B). In addition, similar levels of expression of  $\alpha$ -amylase and human LF mRNA were seen in transformant #1 (Fig.3B and 3C).

#### Example 5

##### Purification of Recombinant human LF

LF was purified from the growth medium using CM Sephadex C50 essentially as described by Stowell, et al., *Biochem J.*, 276:349-59 (1991). The column was preequilibrated with 500 ml of 0.025 M Tris HCl, pH 7.5 0.1M NaCl. The pH of the culture medium was adjusted to pH 7.4 before applying to the preequilibrated column. The column was washed with 500 ml of equilibration buffer and followed by a linear salt gradient from 0.1 to 1.1 M NaCl. Fractions (7 ml total) were assayed for lactoferrin content and purity using SDS/PAGE and silver staining. Fractions containing LF were dialyzed against 0.025 M Tris HCl, pH 7.5/0.1M NaCl and lyophilized.

#### Example 6

##### Quantitation of human LF

Recombinant lactoferrin was quantitated using an ELISA assay essentially as described by Vilja et al., *J. Immunol. Methods*, 76:73-83 (1985). A sensitivity of 5 ng of lactoferrin was obtained using the non-competitive Avidin-biotin assay. Human LF isolated from breast milk (Sigma) was used as standard. Biotinylated human lactoferrin IgG was obtained from Jackson Immunoresearch laboratories, West Grove, PA.

-12-

### Example 7

#### N-terminal Sequencing

Five  $\mu$ g of purified recombinant human LF was resolved on an SDS-polyacrylamide gel and transferred to Problott, a polyvinylidene difluoride-type membrane, following manufacturers instructions (Applied Biosystems). Human LF was detected with Comassie Brilliant Blue staining and destained. This human LF band was excised, washed thoroughly with distilled  $H_2O$  and air-dried. The N-terminal amino acid sequence of the first ten amino acids of human LF was determined by the automated Edman degradation procedure using an applied Biosystems Pulsed-liquid phase sequencer (Model 477A).

With reference to FIGURE 4, panel A illustrates a Silver stained SDS-polyacrylamide gel analysis of recombinant human LF secretion and purification. Lane 1 contains breast milk human LF standard (500 ng). Lanes 2 and 3 contain samples of the growth medium (40  $\mu$ g) from induced control AO7 and transformant #1 respectively. Lanes 4-8 contain 100  $\mu$ l aliquots of eluted fractions (#25, 30, 35, 40, and 45 respectively) collected from the CM-sephadex purification of recombinant LF from the growth medium of transformant #1. The position of the molecular weight markers (BioRad laboratories Richmond, CA) are indicated on the left. Sizes are given in kilodaltons. Panel B illustrates a Western immunoblot analysis of duplicate samples as described in panel A using a specific polyclonal antibody directed against human LF with detection with  $^{125}I$ -protein A. Panel C illustrates #6 N-terminal amino acid sequence of recombinant human LF. Recombinant human LF was sequenced from the N-terminus through 10 residues and is identical to breast milk human LF with the exception of the additional alanine generated in our construction to provide the  $\alpha$ -amylase signal sequence cleavage site.

-13-

Example 8Deglycosylation

Deglycosylation was performed using N-glycosidase F (Boehringer Mannheim). *A. oryzae* growth medium containing 0.5 µg lactoferrin was denatured for 3 minutes at 100°C in the presence of 0.01% SDS. Standard LF from human milk was treated similarly. The samples were subsequently placed on ice for five minutes. N-glycosidase F reactions were conducted in 0.4 M sodium phosphate, (pH 6.8); 0.08% Triton; 0.1% β-mercaptoethanol and 1 unit of enzyme and incubated at 37°C for sixteen hours. PAGE and western analysis was performed using an IgG specifically directed against human lactoferrin to detect an increase in mobility of digested samples.

With reference to FIGURE 5, recombinant human LF was characterized. Panel A illustrates the deglycosylation of lactoferrin. Western analysis of glycosylated and deglycosylated lactoferrin using a specific polyclonal antibody was directed against human lactoferrin with detection with <sup>125</sup>I-protein A. The first panel contains authentic breast milk human LF (500 ng) untreated (-) and treated (+) with N glycosidase F. The second panel contains purified recombinant human LF (500 ng) untreated (-) and treated (+) with N-glycosidase F. The size of glycosylated human LF is indicated with the arrow. Panel B illustrates a functional analysis of recombinant lactoferrin with regard to iron-binding capacity. Panel A and B show the <sup>59</sup>Fe filter binding assay of duplicate samples of authentic breast milk human LF and purified recombinant human LF, respectively, at the concentrations indicated. The first lane in both panels contain BSA (5 µg) as a negative control.

Lactoferrin contains two N-acetylactamine type glycans attached through N-glycosidic linkages. To determine if recombinant lactoferrin was glycosylated correctly, the protein was treated with N-glycosidase F, resolved on SDS-polyacrylamide electrophoresis, transferred to nitrocellulose and probed using a specific IgG directed against human

-14-

lactoferrin (Fig. 5A). N-glycosidase F hydrolyses at the glycosylamine linkage generating a carbohydrate free peptide of smaller molecular weight. Comparison of recombinant LF with purified LF from human milk, illustrates that both proteins co-migrate upon digestion with N-glycosidase F suggesting that the recombinant protein has a glycosylation pattern similar to native LF.

Lactoferrin has a bilobal structure with each lobe having the capacity to bind tightly, but reversibly, one  $\text{Fe}^{3+}$  ion. The iron-binding properties of lactoferrin are crucial for its functional roles. To test if recombinant human LF expressed and secreted in *A. oryzae* has an iron binding capacity similar to authentic lactoferrin, an  $^{59}\text{Fe}$  micro filter binding assay was developed. Purified human lactoferrin isolated from the growth medium of transformant #1 was dialyzed against 0.1M citric acid (pH 2.0) to generate apo-human LF. Native lactoferrin from human milk was treated similarly. Excess  $^{59}\text{Fe}$  (0.2 mCi) was added to these samples in an equal volume of 1 M bicarbonate, followed by incubation at 37°C for 30 minutes. Samples were applied to nitrocellulose membrane and washed several times with bicarbonate. The filter was visualized by autoradiography and Fe-binding was quantitated using a betagon blot analyzer. As illustrated in Fig. 5B, both recombinant and native LF showed a similar level of iron binding at all concentrations tested. The results demonstrate that recombinant human LF is indistinguishable from native human LF in its capacity to bind iron.

With reference to Figure 6, the complete cDNA sequence for human lactoferrin protein is depicted. The cDNA coding for lactoferrin is used to create plasmids and transform eucaryotic cells and to produce the lactoferrin protein.

Strains of aspergillus used in the present invention are auxotrophic mutants that contain a defective *pyr 4* gene that results in an inability to synthesize orotidine 5' phosphate (OMP) decarboxylase. The enzyme is required for uridine synthesis. The strain cannot grow on



-15-

media lacking uridine. The plasmid contains a selectable marker, i.e., a sequence that encodes the gene for OMP decarboxylase. Uptake of the plasmid by the aspergillus can therefore be selected for by growth on media lacking uridine. The aspergillus is transformed by the plasmid such that it can grow on the uridine deficient media.

In one embodiment of the present invention, biologically active recombinant lactoferrin protein is produced. This method comprises synthesizing sequences containing a selectable marker gene, a promoter, a transcription termination sequence and a linker sequence. Subsequently, the sequences are cloned to form a plasmid and the plasmid is digested with a restriction endonuclease. A cDNA coding for lactoferrin is inserted into a restriction site and eucaryotic cells are then transformed with the plasmid expressing the lactoferrin cDNA.

The selectable marker gene useful in the method of the present invention may be any that permits isolation of cells transformed with a lactoferrin cDNA plasmid. Preferably, the selectable marker gene is selected from pyr4, pyrG, argB, trpC and andS.

The promoter useful in the present invention may be any that allows regulation of the transcription of the lactoferrin cDNA. Preferably, the promoter is selected from the group of alcohol dehydrogenase, argB,  $\alpha$ -amylase and glucoamylase.

The transcription termination sequence useful in the present method may be any that allows stabilization of the lactoferrin mRNA. Preferably, the transcription termination sequence is derived from  $\alpha$ -amylase, glucoamylase, alcohol dehydrogenase or benA.

The linker sequence useful in the present method may be any that contains a translation initiation codon, a secretory signal and a restriction enzyme cleavage site. Preferably, the linker element is derived from  $\alpha$ -amylase, glucoamylase or lactoferrin.

The eucaryotic cells useful in the present invention are any that allow for integration of a plasmid comprising the lactoferrin cDNA

-16-

and expression of the lactoferrin cDNA. Preferably, the eucaryotic cells are fungal cells or insect cells. Insect cells such as SF9 are useful in the method of the present invention. More preferably, the fungal cells are yeast cells. Most preferably, the eucaryotic cells useful in the present invention are aspergillus strains, such as *A. oryzae*, *A. Niger*, *A. Nidulans* and *A. Awamori*.

In conclusion, it is seen that the present invention and the embodiments disclosed herein are well adapted to carry out the objectives and obtain the end set forth in this application. Certain changes can be made in the method and apparatus without parting from the spirit and scope of this invention. It is realized that changes are possible and that it is further intended that each element or step presided in any of the filing claims is to be understood as to referring to all equivalent elements or steps for accomplishing the essentially the same results in substantially the same or equivalent manner. It is intended to cover the invention broadly in whatever form its principles may be utilized. The present invention, therefore, is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as others inherent therein.

What is claimed is:

-17-

1. A method for producing biologically active recombinant lactoferrin comprising the steps of:

combining sequences containing a selectable marker gene, a promoter, a transcription termination sequence, and a linker sequence;  
cloning said sequences to form a plasmid;  
digesting said plasmid with a restriction endonuclease;  
inserting a cDNA coding for lactoferrin into a restriction site;  
and  
transforming eucaryotic cells with said plasmid expressing lactoferrin cDNA.

2. The method of Claim 1, wherein said selectable marker gene is selected from the group consisting of pyr4, pyrG, andS, argB and trpC.

3. The method of Claim 1, wherein said cell expresses lactoferrin.

4. Lactoferrin produced by the method of Claim 2.

5. The method of Claim 1, wherein said promoter is selected from the group consisting of alcohol dehydrogenase, argB,  $\alpha$ -amylase, glucoamylase, and benA.

6. The method of Claim 1, wherein said transcription termination sequence is selected from the group consisting of  $\alpha$ -amylase, glucoamylase, alcohol dehydrogenase and benA.

7. The method of Claim 1, wherein said linker sequence is selected from the group consisting of  $\alpha$ -amylase, glucoamylase and lactoferrin.

-18-

8. A plasmid adopted for expression in a eucaryotic cell which comprises the cDNA of Figure 6 and the regulatory elements necessary for the expression of the cDNA in the eucaryotic cell.

9. A plasmid designated pAhLFG.

10. A eucaryotic cell containing the plasmid of Claim 8.

11. The eucaryotic cell of Claim 10, wherein said eucaryotic cell is selected from the group consisting of fungi and insect cells.

12. The eucaryotic cell of Claim 11, wherein said insect cell is SF9.

13. The eucaryotic cell of Claim 11, wherein said fungal cell is yeast.

14. The eucaryotic cell of Claim 13, wherein said yeast cell is *aspergillus*.

15. The eucaryotic cell of Claim 14, wherein said strain of *aspergillus* is selected from the group consisting of *A. oryzae*, *A. Niger*, *A. Nidulans* and *A. Awamori*.

16. A process for producing lactoferrin which comprises culturing a transformant eucaryotic cell containing a recombinant plasmid, said plasmid comprising a plasmid vector having a polydeoxyribonucleotide which codes for lactoferrin proteins in a suitable nutrient medium until lactoferrin protein is formed and, isolating the human lactoferrin.

17. A recombinant expression vector having a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role

-19-

in gene expression; (2) cDNA coding for human lactoferrin; and (3) appropriate transcription and translation initiation and termination sequences.

18. The vector of Claim 17, wherein said genetic element is a promotor.

19. The vector of Claim 18, wherein said promotor is selected from the group consisting of alcohol dehydrogenase, argB,  $\alpha$ -amylase, glucoamylase, and benA.

20. The vector of Claim 17, wherein said transcription termination sequence is selected from the group consisting of  $\alpha$ -amylase, glucoamylase, alcohol dehydrogenase and benA.

21. The protein product of Claim 16.




FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C



FIG. 4A





FIG. 4B



C.

- |    |                             |   |
|----|-----------------------------|---|
|    | α-Amylase Signal Sequence   | Mature α-Amylase                        |
| 1) | Ala Leu Ala                 | Ala Thr Pro Ala Asp Trp Arg Ser Gln Ser |
|    | Lactoferrin Signal Sequence | Mature Human Lactoferrin                |
| 2) | Cys Leu Ala                 | Gly Arg Arg Arg Arg Ser Val Gln Trp Cys |
|    | α-Amylase Signal Sequence   | Mature Recombinant Lactoferrin          |
| 3) | Ala Leu Ala                 | Ala Gly Arg Arg Arg Arg Ser Val Gln Trp |

FIG. 4C




FIG. 5A




FIG. 5B

cDNA SEQUENCE AND AMINO ACID SEQUENCE  
OF HUMAN LACTOFERRIN

1  
GAATTCC GACCGCAGAC

18  
ATG AAA CTT GTC TTC CTC GTC CTG CTG TTC CTC GGG GCC CTC GGA CTG  
met lys leu val phe leu val leu leu phe leu gly ala leu gly leu  
1

66  
TGT CTG GCT GGC CGT AGG AGA AGG AGT GTT CAG TGG TGC ACC GTA TCC  
cys leu ala gly arg arg arg arg ser val gln trp cys thr val ser  
17

114  
CAA CCC GAG GCC ACA AAA TGC TTC CAA TGG CAA AGG AAT ATG AGA AGA  
gln pro glu ala thr lys cys phe gln trp gln arg asn met arg arg  
33

162  
GTG CGT GGC CCT CCT GTC AGC TGC ATA AAG AGA GAC TCC CCC ATC CAG  
val arg gly pro pro val ser cys ile lys arg asp ser pro ile gln  
49

210  
TGT ATC CAG GCC ATT GCG GAA AAC AGG GCC GAT GCT GTG ACC CTT GAT  
cys ile gln ala ile ala glu asn arg ala asp ala val thr leu asp  
65

258  
GGT GCT TTC ATA TAC GAG GCA GGC CTG GCC CCC TAC AAA CTG CGA CCT  
gly gly phe ile tyr glu ala gly leu ala pro tyr lys leu arg pro  
81

306  
GTA GCG GCG GAA GTC TAC GGG ACC GAA AGA CAG CGA CGA ACT CAC TAT  
val ala ala glu val tyr gly thr glu arg gln pro arg thr his tyr  
97

354  
TAT GCC GTG GCT GTG GTG AAG AAG GGC GGC AGC TTT CAG CTG AAC GAA  
tyr ala val ala val val lys lys gly gly ser phe gln leu asn glu  
113

402  
CTG CAA GGT CTG AAG TCC TGC CAC ACA GGC CTT CGC AGC ACC GCT CGA  
leu gln gly leu lys ser cys his thr gly leu arg arg thr ala gly  
129

450  
TGC AAT CTG CCT ATA GCG ACA CTT CGT CCA TTC TTG AAT TGG AGC GGT  
trp asn val pro ile gly thr leu arg pro phe leu asn trp thr gly  
145

Sheet 1 of 4

FIGURE 6

498  
CCA CCT GAG CCC ATT GAG GCA GCT GTG GCC AGG TTC TTC TCA GCC AGC  
pro pro glu pro ile glu ala ala val ala arg phe phe ser ala ser  
161

546  
TGT GTT CCC GGT GCA GAT AAA GGA CAG TTC CCC AAC CTG TGT CGC CTG  
cys val pro gly ala asp lys gly gln phe pro asn leu cys arg leu  
177

594  
TGT GCG GGG ACA GGG GAA AAC AAA TGT GCC TTC TCC TCC CAG GAA CCG  
cys ala gly thr gly glu asn lys cys ala phe ser ser gln glu pro  
193

642  
TAC TTC AGC TAC TCT GCT GCC TTC AAG TGT CTG AGA GAC GGC GCT GGA  
tyr phe ser tyr ser gly ala phe lys cys leu arg asp gly ala gly  
209

690  
GAC GTG GCT TTT ATC ACA GAG ACC ACA GTG TTT GAG GAC CTG TCA GAC  
asp val ala phe ile arg glu ser thr val phe glu asp leu ser asp  
225

738  
GAG GCT GAA AGG GAC GAG TAT GAG TTA CTC TGC CCA GAC AAC ACT CGG  
glu ala glu arg asp glu tyr glu leu leu cys pro asp asn thr arg  
241

786  
AAG CCA GTG GAC AAG TTC AAA GAC TGC CAT CTG GCC CGG GTC CCT TCT  
lys pro val asp lys phe lys asp cys his leu ala arg val pro ser  
257

834  
CAT GCC GTT GTG GCA CGA AGT GTC AAT GGC AAG GAG GAT GCC ATC TGG  
his ala val val ala arg ser val asn gly lys glu asp ala ile trp  
273

882  
AAT CTT CTC CGC CAG GCA CAG GAA AAG TTT GGA AAG GAC AAG TCA CCG  
asn leu leu arg gln ala gln gln lys phe gly lys asp lys ser pro  
289

930  
AAA TTC CAG CTC TTT GGC TCC CCT AGT GGG CAC AAA GAT CTG CTG TTC  
lys phe gln leu phe gly ser pro ser gly gln lys asp leu leu phe  
305

978  
AAG GAC TCT GCC ATT GCG TTT TCG AGC GTG CCC CCG AGG ATA GAT TCT  
lys asp ser ala ile gly phe ser arg val pro pro arg ile asp ser  
321

1026  
GGG CTG TAC CTT GGC TCC GGC TAC TTC ACT GCC ATC CAG AAC TTG AGG  
gly leu tyr leu gly ser gly tyr phe thr ala ile gln asn leu arg  
337

1074  
AAA AGT CAG GAG GAA GTG GCT GCC CGG CGT GCG CGG GTC GTG TGG TGT  
lys ser glu glu glu val ala ala arg arg ala arg val val trp cys  
353

1122  
GCG GTG GGC GAG CAG GAG CTG GCG AAG TGT AAC CAG TGG AGT GGC TTC  
ala val gly glu gln glu leu arg lys cys asn gln trp ser gly leu  
369

1170  
AGC GAA GGC AGC GTG ACC TGC TCC TCG GCC TCC ACC ACA GAG GAC TGC  
ser glu gly ser val thr cys ser ser ala ser thr thr glu asp cys  
385

1218  
ATC GCC CTG GTG CTG AAA GGA GAA GCT GAT GCC ATG AGT TTG GAT GGA  
ile ala leu val leu lys gly glu ala asp ala met ser leu asp gly  
401

1266  
GGA TAT GTG TAC ACT GCA GGC AAA TGT GGT TTG GTG CCT GTC CTG GCA  
gly tyr val tyr thr ala gly lys cys gly leu val pro val leu ala  
417

1314  
GAG AAC TAC AAA TCC CAA CAA AGC AGT GAC CCT GAT CCT AAC TGT GTG  
glu asn tyr lys ser gln gln ser ser asp pro asp pro asn cys val  
433

1362  
GAT AGA CCT GTG GAA GGA TAT CTT GCT GTG GCG GTG GTT ACC AGA TCA  
asp arg pro val glu gly tyr leu ala val ala val val arg arg ser  
449

1410  
GAC ACT AGC CTT ACC TGG AAC TCT GTG AAA GGC AAC AAG TCC TGC CAC  
asp thr ser leu thr trp asn ser val lys gly lys lys ser cys his  
465

1458  
ACC GCC GTG GAC AGG ACT GCA GGC TGG AAT ATC CCC ATG GGC CTG CTC  
thr ala val asp arg thr ala gly trp asn ile pro met gly leu leu  
481

1506  
TTC AAC CAG ACG GGC TCC TGC AAA TTT GAT GAA TAT TTC AGT CAA AGC  
phe asn gln thr gly ser cys lys phe asp glu tyr phe ser gln ser  
497

1554  
TGT GCC CCT GGS TCT GAC CCG AGA TCT AAT CTC TGT GCT CTG TGT ATT  
cys ala pro gly ser asp pro arg ser asn leu cys ala leu cys ile  
513

1602  
GCG GAC GAG CAG GGT GAG AAT AAG TGC GTG CCC AAC AGC AAT GAG AGA  
gly asp glu gln gly glu asn lys cys val pro asn ser asn glu arg  
529

1650  
TAC TAC GGC TAC ACT GGG GCT TTC CGG TGC CTG GCT GAC AAT GCT GGA  
tyr tyr gly tyr thr gly ala phe arg cys leu ala glu asn ala gly  
545

1698  
GAC GTT GCA TTT GTG AAA GAT GTC ACT GTC TTG CAG AAC ACT GAT GGA  
asp val ala phe val lys asp val thr val leu gln asn thr asp gly  
561

1746  
AAT AAC AAT GAG GCA TGG GCT AAG GAT TTG AAG CTG GCA GAC TTT GCG  
asn asn asn glu ala trp ala lys asp leu lys leu ala asp phe ala  
577

1794  
CTG CTG TGC CTC GAT GCC AAA CGG AAG CCT GTG ACT GAG GCT AGA AGC  
leu leu cys leu asp gly lys arg lys pro val thr glu ala arg ser  
593

1842  
TSC CAT CTT GCC ATG GCC CCG AAT CAT GCC GTG GTG TCT CGG ATG GAT  
cys his leu ala met ala pro asn his ala val val ser arg met asp  
609

1890  
AAG GTG GAA CGC CTG AAA CAG GTG CTG CTC CAC CAA CAG GCT AAA TTT  
lys val glu arg leu lys gln val leu leu his gln gln ala lys phe  
625

1938  
GGG AGA AAT GGA TCT GAC TGC CCG GAC AAG TTT TGC TTA TTC CAG TCT  
gly arg asn gly ser asp cys pro asp lys phe cys leu phe gln ser  
641

1986  
GAA ACC AAA AAC CTT CTG TTC AAT GAC AAC ACT GAG TGT CTG GCC AGA  
glu thr lys asn leu leu phe asn asp asn thr glu cys leu ala arg  
657

2034  
CTC CAT GGC AAA ACA ACA TAT GAA AAA TAT TTG GGA CCA CAG TAT GTC  
leu his gly lys thr thr tyr glu lys tyr leu gly pro gln tyr val  
673

2082  
GCA GGC ATT ACT AAT CTG AAA AAG TGC TCA ACC TCC CCC CTC CTG GAA  
ala gly ile thr asn leu lys lys cys ser thr ser pro leu leu glu  
689

2130  
GCC TGT GAA TTC CTC AGG AAG TAA  
ala cys glu phe leu arg lys \*\*\* ACCGAA GAAGATGCCC CAGCTCCCA  
705

2180  
AGAAAGGCTC AGCCATTGAC TGCCCCAGC TCTTCTGCCC AGGTGTGTG GGGCCTTGGC

2240  
TCCCCCTCTG AAGGTGGGA TTGCCATCC ATCTGCTTAC AATTCCCTGC TGTCGTCTTA

2300  
GCAAGAAGTA AATGAGAAA TTTTGTGAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/03614

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : C07K 15/06, 15/14; C12N 15/04, 15/06, 15/10, 15/12  
 US CL : 536/23.5; 530/350, 394; 435/69.1, 240.2, 254, 255, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.5; 530/350, 394; 435/69.1, 240.2, 254, 255, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, DIALOG, Swissprot, PIR, search terms: lactoferrin, amylase, oryzae, aspergillus, cDNA

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	Biochemical Journal, Volume 276, issued June 1991, Stowell et al., "Expression of cloned human lactoferrin in baby-hamster kidney cells", pages 349-355, see especially p. 350, column 2.	<u>1, 3, 4, 8, 10, 11, 13</u> <u>16-18, 21</u> 1-21
Y	WO, A, 89/01969 (Woldike, H.F.) 09 March 1989, see whole publication, especially pages 3, 5, 6, 10 and 11.	1-21
Y	Gene, Volume 79, issued 1989, Gines et al., "Aspergillus oryzae has two nearly identical Taka-amylase genes, each containing eight introns", pages 107-117, see especially the abstract and Figures 2 and 3.	1-21
Y	Journal of Molecular Biology, Volume 173, issued 1984, G. von Heijne, "How signal sequences maintain cleavage specificity", pages 243-251, see especially page 244.	1-21

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	*+ later document published after the international filing date or priority date and not in conflict with the application but cited to substantiate the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claims or which is cited to establish the publication date of another citation or other special reason (as specified)	
*O* document referring to no oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	*A* document member of the same patent family

Date of the actual completion of the international search

27 May 1993

Date of mailing of the international search report

27 JUL 1993

Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT  
 Washington, D.C. 20231

Authorized officer

KEITH C. FURMAN, PH.D.

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1992)\*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US93/03614

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Nucleic Acids Research, Volume 18, Number 13, issued 1990, Powell et al., "Nucleotide sequence of human lactoferrin cDNA", page 4013.	1-21
Y	Lipids, Volume 24, Number 9, issued 1989, Høge-Jensen et al., "Rhizomucor miehei triglyceride lipase is processed and secreted from transformed <u>Aspergillus oryzae</u> ", pages 781-785, see especially the abstract.	1-21
Y	US, A, 4,740,461 (Kaufman) 26 April 1988, see whole patent, especially Table 1 and columns 11 and 12.	1-21
Y,P	US, A, 5,155,037 (Summers) 13 October 1992, filed 04 August 1989, see whole patent, especially columns 1 and 4-8.	1, 3, 8, 10, 11, 12, 16-18, 21